

# The Brunn-Minkowski Inequality and nontrivial cycles in the discrete torus

Noga Alon <sup>\*</sup>      Ohad N. Feldheim <sup>†</sup>

March 21, 2010

## Abstract

Let  $(C_m^d)_\infty$  denote the graph whose set of vertices is  $Z_m^d$  in which two distinct vertices are adjacent iff in each coordinate they are either equal or differ, modulo  $m$ , by at most 1. Bollobás, Kindler, Leader and O'Donnell proved that the minimum possible cardinality of a set of vertices of  $(C_m^d)_\infty$  whose deletion destroys all topologically nontrivial cycles is  $m^d - (m-1)^d$ . We present a short proof of this result, using the Brunn-Minkowski Inequality, and also show that the bound can only be achieved by selecting a value  $x_i$  in each coordinate  $i$ ,  $1 \leq i \leq d$ , and by keeping only the vertices whose  $i$ -th coordinate is not  $x_i$ , for all  $i$ .

**Keywords:** Brunn-Minkowski, Discrete torus, Nontrivial cycles.

**AMS classification:** 05C35,05C38,05C40.

## 1 Introduction

Let  $(C_m^d)_\infty$  denote the graph whose set of vertices is  $Z_m^d$  in which two distinct vertices are adjacent iff in each coordinate they are either equal or differ, modulo  $m$ , by at most 1. This graph is the product of  $d$  copies of the cycle of length  $m$ , and can be viewed as the graph of the discrete torus. The problem of determining the minimum possible cardinality of a set of vertices of this graph that intersects all noncontractible cycles in it, has been considered by Saks, Samorodnitsky, and Zosin in [1], motivated by the problem of exhibiting directed multi-commodity problems that have a large integrality gap. Their estimate has been improved to a tight one, which is  $m^d - (m-1)^d$ , by Bollobás, Kindler, Leader and O'Donnell in [2], where a connection to the parallel repetition of the odd cycle game is mentioned. In this note we describe a short intuitive proof of the same result, based on

---

<sup>\*</sup>Schools of Mathematics and Computer Science, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel. Email: nogaa@tau.ac.il. Research supported in part by an ERC Advanced grant, by a USA-Israeli BSF grant, and by the Hermann Minkowski Minerva Center for Geometry in Tel Aviv University.

<sup>†</sup>School of Mathematics, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel. Email: ohad.f@netvision.net.il. Research supported in part by an ERC Advanced grant.

the Brunn-Minkowski isoperimetric Inequality. The proof also implies that equality is achieved only when the  $(m - 1)^d$  vertices remaining form the graph of a  $d$ -dimensional hypercube of edge length  $m - 1$ , that is, the product of  $d$  paths, each having  $m - 1$  vertices.

It is worth noting that the problem of determining the minimum cardinality of a set of *edges* of the graph  $(C_m^d)_\infty$  that intersects all nontrivial cycles, discussed in [3], [4], seems more difficult and only an asymptotic estimate of this minimum is known.

## 2 The proof

Let  $Z_m^d$  be the set of vertices of  $(C_m^d)_\infty$ , and consider them as points in  $\mathbb{Z}^d$ . It is convenient to view  $\mathbb{Z}^d$  as an infinite graph in which two distinct vectors are adjacent iff they differ in at most 1 in each coordinate. For two vectors  $\bar{a} = (a_1, a_2, \dots, a_d)$  and  $\bar{b} = (b_1, b_2, \dots, b_d)$  in  $Z_m^d$  or in  $\mathbb{Z}^d$  we write that  $\bar{b} \nearrow \bar{a}$  iff  $a_i - b_i \in \{0, 1\}$  for all  $i$ . Note that  $\nearrow$  is a reflexive relation. Note also that the following holds:

**Observation 1.** *If  $\bar{b}_1, \bar{b}_2 \nearrow \bar{a}$ , then  $\bar{b}_1$  and  $\bar{b}_2$ , considered as vertices of  $(C_m^d)_\infty$ , are either equal or connected.*

Recall that the Brunn-Minkowski Inequality, generalized by Lusternik (see, e.g. [5]), is the following.

**Theorem** (The Brunn-Minkowski Inequality). *Let  $n \geq 1$  and let  $\mu$  be the Lebesgue measure on  $\mathbb{R}^n$ . Define  $A + B := \{a + b \in \mathbb{R}^n \mid a \in A, b \in B\}$ . Let  $A$  and  $B$  be two nonempty compact subsets of  $\mathbb{R}^n$ . The following inequality holds:*

$$[\mu(A + B)]^{1/n} \geq [\mu(A)]^{1/n} + [\mu(B)]^{1/n}.$$

*Equality is achieved iff  $A$  and  $B$  are homothetic (that is, one is a rescaled version of the other)*

Using Brunn-Minkowski we obtain the following useful lemma:

**Lemma 2.1.** *Let  $S \subseteq \mathbb{Z}^d$ . Suppose  $S^+ = \{\bar{a} \mid \exists \bar{b} \in S (\bar{b} \nearrow \bar{a})\}$ , then  $\sqrt[d]{|S^+|} \geq \sqrt[d]{|S|} + 1$ , and equality holds iff  $S$  is a hypercube.*

*Proof.* define  $\widehat{S} = \bigcup_{\bar{a} \in S} \{\prod_{i \in \{1, \dots, d\}} [a_i - 1, a_i]\}$ , and note that  $|S| = \mu(\widehat{S})$ . It is easy to check that  $\widehat{S}^+ = \widehat{S} + [0, 1)^d$ . Plugging this and the fact that  $|S^+| = \mu(\widehat{S}^+)$  into the Brunn-Minkowski Inequality, the result follows.  $\square$

We can now state and prove the main theorem:

**Theorem 1.** *If  $S \subset Z_m^d$  is a set of vertices of  $Z_m^d$  that does not contain any non-contractible cycle of the torus, then  $|S| \leq (m - 1)^d$ . Equality holds if and only if  $S$  is a hypercube with edges of size  $m - 1$ .*

*Proof.* Striving for contradiction, suppose that either  $|S| > (m-1)^d$ , or  $|S| = (m-1)^d$  but  $S$  is not a hypercube. Denote the connected components of  $S$  by  $C_1, \dots, C_k$ . Pick a vertex representative for each component  $C_i$ , and denote it by  $\bar{c}_i$ . Let the natural projection from  $\mathbb{Z}^d$  into  $Z_m^d$  be  $\pi(\bar{x})$ . Slightly abusing notation, denote by  $\pi^{-1}(C_i)$  the connected component of  $\bar{c}_i$  in  $\pi^{-1}(S)$ , regarding here  $\bar{c}_i$  as an element of  $\mathbb{Z}^d$ . (This is instead of taking the whole  $\pi$  pre-image of  $C_i$ ). As  $S$  contains no non-trivial cycle,  $\pi^{-1}(C_i)$  must be finite for all  $i$ . We next show that there exist two distinct preimages of some vertex  $\bar{a}$  in one of the connected components  $C_i$  of  $S$ , implying that it contains a nontrivial cycle, and thus contradicting the assumption.

Define  $\tilde{S} = \bigcup_{i=1}^k \pi^{-1}(C_i)$ . Since every vertex in  $S$  has a unique corresponding vertex in  $\tilde{S}$  we deduce that  $|S| = |\tilde{S}|$ . Looking at  $\tilde{S}^+ = \{\bar{a} | \exists \bar{b} \in \tilde{S} (\bar{b} \nearrow \bar{a})\}$  we can apply our assumption and lemma 2 to conclude that  $|\tilde{S}^+| > m^d$ . By The Pigeonhole Principle we deduce the existence of  $\bar{a}_1 \neq \bar{a}_2$  in  $\tilde{S}^+$  such that  $\pi(\bar{a}_1) = \pi(\bar{a}_2)$ . By the definition of  $\tilde{S}^+$  there must be two elements  $\bar{b}_1, \bar{b}_2 \in \tilde{S}$  such that  $\bar{b}_1 \nearrow \bar{a}_1$  and  $\bar{b}_2 \nearrow \bar{a}_2$ . By Observation 1 we know that  $\pi(\bar{b}_1)$  and  $\pi(\bar{b}_2)$  are connected in  $S$  and thus  $\bar{b}_1$  and  $\bar{b}_2$  belong to the same connected component  $\pi^{-1}(C_i)$  of  $\tilde{S}$ , for some  $i$ . Denote  $\bar{b}'_1 = \bar{a}_2 - \bar{a}_1 + \bar{b}_1$ . Note that  $\bar{b}'_1 \neq \bar{b}_1$ ,  $\pi(\bar{b}'_1) = \pi(\bar{b}_1)$ , and  $\bar{b}'_1 \nearrow \bar{a}_2$ , since  $\bar{a}_2 - \bar{b}'_1 = \bar{a}_2 - (\bar{a}_2 - \bar{a}_1 + \bar{b}_1) = \bar{a}_1 - \bar{b}_1$ .

By Observation 1 we conclude that  $\bar{b}'_1$  and  $\bar{b}_2$  are either equal or connected. As  $\bar{b}_2 \in \pi^{-1}(C_i)$  we conclude that  $\bar{b}'_1 \in \pi^{-1}(C_i)$ , which leads to contradiction, since  $\bar{b}_1$  also lies in  $C_i$ . Therefore, either  $|S| = (m-1)^d$  and  $S$  is a hypercube, or  $|S| < (m-1)^d$ , completing the proof.  $\square$

## References

- [1] N. Alon and B. Klartag, Economical toric spines via Cheeger's Inequality, *Journal of Topology and Analysis* 1 (2009), 101–111.
- [2] Bollobás, Kindler, Leader and O'Donnell, Eliminating cycles in the discrete torus, *Algorithmica* 50 (2007), pp. 446–454.
- [3] R. Raz, A counterexample to strong parallel repetition, *Proc. FOCS 2008*, pp. 369–373.
- [4] M. E. Saks, A. Samorodnitsky and L. Zosin, A lower bound on the integrality gap for minimum multicut in directed networks, *Combinatorica* 24 (2004), pp. 525–530.
- [5] R. Schneider, *Convex Bodies: the Brunn-Minkowski Theory*, Cambridge University Press, Cambridge, (1993).